## REPORT OF EXAMINATION OF ALASKA SALMON STREAM

## DEPARTMENT OF COMMERCE BUREAU OF FISHERIES



Name of stream at head of Kenalku Bay Date examined

Observed during season 1930.

Location Kenalku Bay

Approximate length 1-1/4 miles

Width

35 ft

Depth inches

Condition and stage (whether clear, high, low, etc.) ATTRAGE.

Lakes, if any, size, character, and whether accessible from sea Solid rock falls 1/2 mile from high water, sockeyes and cohoes go over these falls but humpies go only to foot of falls.

Salmon run (heavy, light, average, etc.) Average.

Date and duration of run of each species

Sockeye run from July 1 to July 25,

Humpbacks Aug. 5 to Sept. 5, Peak run Aug. 10 to 25

Cohoes run later.

Estimated number of each kind of salmon on spawning beds, or in stream

With the exception of 296 sockeyes taken in the bay, all fish went to spawning grounds. No humpbacks taken.

Former size of runs of each species and evidence of depletion, if any

About the same as the past several seasons up to closed season.

Natural barriers and whether removable

Solid rock falls.

Where markers are placed

1500 feet from Mouth of stream on trees.

Remarks, sketches, etc.

SEE Salmon school mostly at mouth of stream however they go back of dead line at times.

Harry Peehan, Stream Watchman.

(Name and title of employee making examination)

Send original to Washington office, one copy to Seattle office, and one copy to Juneau office. Use reverse side if additional space is required, or explanatory sketches are made which can not go on this side. U. S. GOVERNMENT PRINTING OFFICE: 1929